



How is this measured?

Sources and definitions for quality measures reported on Washington HealthCareCompare

Measures groups included on the website:

- Quality measures for medical practices
- Quality measures for hospitals
- Quality measures for Healthier Washington (Accountable Communities for Health)
- Quality measures by insurance market sectors

WA-APCD MEDICAL PRACTICE QUALITY MEASURES

Measure Name	Steward	Reporting Unit	Definition
Medical Practice Rating	WA-APCD	Summary	Global 1–3 star rating
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	NCQA	Percentage	The percentage of adults, ages 18–64 years, with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription.
Ambulatory Care – Avoidable Emergency Department Visits	OR-QCORP	Rate per 1,000 Member Years	The rate of patient ED visits for conditions that could have been more appropriately managed by or referred to a primary care provider.
Antidepressant Medication Management: Effective Acute Phase Treatment	NCQA	Percentage	The percentage of members, ages 18 years and older, who were treated with antidepressant medication, had a diagnosis of major depression, and who remained on an antidepressant medication treatment.
Breast Cancer Screening	NCQA	Percentage	The percentage of women, ages 50–74 years, who had a mammogram to screen for breast cancer.
Cervical Cancer Screening	NCQA	Percentage	The percentage of women, ages 21–64 years, who were screened for cervical cancer using either of two methods defined by the measure (interval dependent upon screening method).
Comprehensive Diabetes Care – Eye Exam Performed	NCQA	Percentage	The percentage of members, ages 18–75 years, with diabetes (Type 1 and/or Type 2) who had an eye exam during the measurement year or the year prior.

Measure Name	Steward	Reporting Unit	Definition
Comprehensive Diabetes Care – HbA1c Testing	NCQA	Percentage	The percentage of members, ages 18–75 years, with diabetes (Type 1 and/or Type 2) who had an HbA1c test during the measurement year.
Comprehensive Diabetes Care – Medical Attention for Nephrology	NCQA	Percentage	The percentage of members, ages 18–75 years, with diabetes (Type 1 and/or Type 2) who received a nephropathy screening or monitoring test or had evidence of nephropathy during the measurement year or the year prior.
Chlamydia Screening in Women	NCQA	Percentage	The percentage of women, ages 16–24 years, who were identified as sexually active and who had at least one test for chlamydia during the measurement year.
Appropriate Testing for Children With Pharyngitis	NCQA	Percentage	The percentage of children, ages 3–18 years, who were diagnosed with pharyngitis, dispensed an antibiotic, and received a group A streptococcus (strep) test for the episode. A higher rate represents better performance (i.e., appropriate testing).
Use of Imaging Studies for Low Back Pain	NCQA	Percentage	The percentage of members, ages 18–50 years, with a primary diagnosis of low back pain who did not have an imaging study (i.e., plain X-ray, MRI, or CT scan) within 28 days of the diagnosis. (Note that a higher score indicates better performance.)
Mental Health Service Penetration	WA DSHS	Percentage	The percentage of members with a mental health service need who received mental health services in the measurement year. Separate reporting for age groups: 6–17 years and 18–64 years.
Annual Monitoring for Patients on Persistent Medications – ACE Inhibitors or ARBs	NCQA	Percentage	The percentage of patients, ages 18 years and older, who received at least 180 treatment days of ambulatory medication therapy for ACE/ARB during the measurement year and had at least one therapeutic monitoring event for the therapeutic agent in the measurement year.
Proportion of Days Covered – Diabetes All Class (DR)	PQA	Percentage	The percentage of patients, ages 18 years and older, who met the proportion of days covered (PDC) threshold of 80% for diabetes roll-up medications.
Proportion of Days Covered – Renin Angiotensin System Antagonists (RASA)	PQA	Percentage	The percentage of patients, ages 18 years and older, who met the proportion of days covered (PDC) threshold of 80% for Renin Angiotensin System Antagonists.
Proportion of Days Covered – Statins (STA)	PQA	Percentage	The percentage of patients, ages 18 years and older, who met the proportion of days covered (PDC) threshold of 80% for statins.
Chronic Obstructive Pulmonary Disease (COPD) or Asthma in Older Adults Admission Rate	AHRQ	Rate per 1,000 Member Years	The rate of adults, ages 40 years and older, who had an admission for a principal diagnosis of chronic obstructive pulmonary disease (COPD) or asthma.

Measure Name	Steward	Reporting Unit	Definition
Well-Child Visits in the First 15 Months of Life	NCQA	Percentage	The percentage of members who turned 15 months old during the measurement year and who had six or more well-child visits with a PCP during their first 15 months of life.
Well-Child Visits in the Third, Fourth, Fifth, and Sixth Years of Life	NCQA	Percentage	The percentage of members, ages 3–6 years, who had one or more well-child visits with a PCP during the measurement year.

WA-APCD HOSPITAL QUALITY MEASURES

Measure Name	Measure Source	Reporting Unit	Description
Hospital Summary Rating	CMS Hospital Compare	Summary	Global 1–5 star rating
30-day All-Cause Hospital Readmissions (HWR)	CMS Hospital Compare	Observed-to-Expected	All-condition risk-adjusted unplanned readmissions within 30 days of discharge
Hospital 30-day Death Rate for Heart attack	CMS Hospital Compare	Per 100 Persons	30-day risk-standardized mortality rate for a hospital. Mortality is defined as death for any cause within 30 days after the date of admission for the index admission, for patients 18 and older discharged from the hospital with a principal diagnosis of acute myocardial infarction.
Medicines Explained When Leaving the Hospital	CMS Hospital Compare	Percentage	HCAHPS patient experience survey: communication about medicines and discharge information
Information When Leaving the Hospital	CMS Hospital Compare	Percentage	HCAHPS patient experience survey: communication about medicines and discharge information
Stroke -Thrombolytic Therapy	CMS Hospital Compare	Percentage	Acute ischemic stroke patients who arrive at hospital within 2 hours of time last known well for whom IV t-PA was initiated at this hospital within 3 hours of time last known well
Patient Safety for 11 Indicators (composite)	CMS Hospital Compare	Observed-to-Expected	Weighted average of the adjusted observed-to-expected ratios for the following component indicators: (1) pressure ulcer rate, (2) iatrogenic pneumothorax rate, (3) postoperative hip fracture rate, (4) postoperative hemorrhage or hematoma, (5) physiologic and metabolic derangement, (6) postoperative respiratory failure, (7) perioperative pulmonary embolism or deep vein thrombosis rate, (8) postoperative sepsis rate, (9) postoperative wound dehiscence rate, and (10) accidental puncture or laceration rate

Measure Name	Measure Source	Reporting Unit	Description
Catheter-associated Urinary Tract Infection - In ICU	WA State Hospital Association	Per 1,000 Persons	Standardized Infection Ratio (SIR) of healthcare-associated, catheter-associated urinary tract infections
Catheter-associated Urinary Tract Infection - Outside ICU	WA State Hospital Association	Per 1,000 Persons	Standardized Infection Ratio (SIR) of healthcare-associated, catheter-associated urinary tract infections
Falls with Injury (per 1,000)	WA State Hospital Association	Per 1,000 Persons	Patient falls of injury level minor or greater by eligible hospital unit
Cesarean Deliveries	WA State Hospital Association	Percentage	Number of nulliparous women with a term, singleton baby in a vertex position delivered by cesarean birth

WA-APCD HEALTHIER WASHINGTON QUALITY MEASURES

Measure Name	Source	Reporting Unit	Population	Description
Emergency Room Visits Per 1,000 Persons	Claims	Visits Per 1,000 Person Months	Medicaid	Number of emergency department visits per 1,000 population and is calculated in member months for Medicaid
Good Mental, Emotional Health	DOH	Percentage	All	Percentage of adults ages 18 and older who answer "14 or more days" in response to the question, "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good."
Avoid Return to Hospital for Mental Health Condition	DSHS	Ratio Observed to Expected	Medicaid	For members 18 years of age and older, the number of acute inpatient psychiatric stays that were followed by an acute readmission for a psychiatric diagnosis within 30 days. Report for Medicaid only.
Adult Tobacco Use	DOH	Percentage	All	The percentage of adults 18 years and older who answer "every day" or "some days" in response to the survey question, "Do you now smoke cigarettes every day, some days, or not at all?"
Unintended Pregnancies	DOH	Percentage	All	Percentage of pregnancies that was unintended at the time of conception

Measure Name	Source	Reporting Unit	Population	Description
Testing for Cause of Low Back Pain	Claims	Percentage	Medicaid	The percentage of members, ages 18–50 years, with a primary diagnosis of low back pain who did not have an imaging study (i.e., plain X-ray, MRI, or CT scan) within 28 days of the diagnosis. (Note that a higher score indicates better performance.)
Avoid Wrong Medication for Bronchitis	Claims	Percentage	Medicaid	The percentage of adults, ages 18–64 years, with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription.
Checking Patients Who Need Ongoing Medications	Claims	Percentage	Medicaid	The percentage of patients, ages 18 years and older, who received at least 180 treatment days of ambulatory medication therapy for ACE/ARB during the measurement year and had at least one therapeutic monitoring event for the therapeutic agent in the measurement year.
Treating Children with Throat Infections	Claims	Percentage	Medicaid	The percentage of children, ages 3–18 years, who were diagnosed with pharyngitis, dispensed an antibiotic, and received a group A streptococcus (strep) test for the episode. A higher rate represents better performance (i.e., appropriate testing).
Avoid Unnecessary Emergency Visits	Claims	Rate per 1,000 Member Years	All	The rate of patient ED visits for conditions that could have been more appropriately managed by or referred to a primary care provider.
Blood Sugar Tested	Claims	Percentage	Medicaid	The percentage of members, ages 18–75 years, with diabetes (Type 1 and/or Type 2) who had an HbA1c test during the measurement year.
Check for Kidney Disease	Claims	Percentage	Medicaid	The percentage of members, ages 18–75 years, with diabetes (Type 1 and/or Type 2) who received a nephropathy screening or monitoring test or had evidence of nephropathy during the measurement year or the year prior.
Eye Exam to Check for Vision Loss	Claims	Percentage	Medicaid	The percentage of members, ages 18–75 years, with diabetes (Type 1 and/or Type 2) who had an eye exam during the measurement year or the year prior.
Getting Medications to Control Blood Sugar	Claims	Percentage	Medicaid	The percentage of patients, ages 18 years and older, who met the proportion of days covered (PDC)

Measure Name	Source	Reporting Unit	Population	Description
				threshold of 80% for diabetes roll-up medications.
Getting Anti-depressant Medications	Claims	Percentage	Medicaid	The percentage of members 18 years and older who were treated with antidepressant medication, had a diagnosis of major depression and who remained on an antidepressant medication treatment: effective acute phase treatment rate
Getting Care for Attention Deficit/Hyperactivity Disorder	Claims	Percentage	Medicaid	The percentage of children newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed. Continuation and Maintenance Phase rate
Follow-Up Visit After Hospital Stay for Mental Health	Claims	Percentage	Medicaid	The percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness diagnoses and who had a follow-up visit with a mental health practitioner within 30 days of discharge
Testing for Lung Disease	Claims	Percentage	Medicaid	The percentage of members 40 years of age and older with a new diagnosis of COPD or newly active COPD, who received appropriate spirometry testing to confirm the diagnosis.
Getting Medications for Asthma	Claims	Percentage	Medicaid	The percentage of members 5-85 years of age who were identified as having persistent asthma and were dispensed appropriate medications that they remained on for at least 75% of their treatment period. (For Medicaid, this measure is reported for people ages 5-64.)
Fewer Hospital Stays Due to Good Lung Care	AHRQ	Rate per 1,000 Member Years	Medicaid	The rate of adults, ages 40 years and older, who had an admission for a principal diagnosis of chronic obstructive pulmonary disease (COPD) or asthma.

Measure Name	Source	Reporting Unit	Population	Description
Getting Medications to Lower Blood Pressure	Claims	Percentage	Medicaid	The percentage of patients, ages 18 years and older, who met the proportion of days covered (PDC) threshold of 80% for Renin Angiotensin System Antagonists.
Getting Medications to Lower Cholesterol	Claims	Percentage	Medicaid	The percentage of patients, ages 18 years and older, who met the proportion of days covered (PDC) threshold of 80% for statins
Infants Get Regular Check-ups	Claims	Percentage	Medicaid	The percentage of members who turned 15 months old during the measurement year and who had six or more well-child visits with a PCP during their first 15 months of life.
Children Get Regular Check-ups	Claims	Percentage	Medicaid	The percentage of members, ages 3-6 years, who had one or more well-child visits with a PCP during the measurement year.
Having a Primary Care Visit Yearly	Claims	Percentage	Medicaid	The percentage of members 12 months - 19 years of age who had a visit with a PCP, all children ages.
Getting Substance Abuse Services	DSHS	Percentage	Medicaid	The percentage of members with a substance use disorder treatment need who received a substance use disorder treatment in the measurement year, age 12-17 years
Getting Mental Health Services	DSHS	Percentage	Medicaid	The percentage of members with a mental health service need who received mental health services in the measurement year age 6-17 years
Having a Doctor/Office Visit Yearly	Claims	Percentage	Medicaid	The percentage of members 20 years and older who had an ambulatory or preventive care visit, all adult ages.
Getting Substance Abuse Services	DSHS	Percentage	Medicaid	The percentage of members with a substance use disorder treatment need who received a substance use disorder treatment in the measurement year, age 18 years and older
Getting Mental Health Services	DSHS	Percentage	Medicaid	The percentage of members with a mental health service need who received mental health services in the

Measure Name	Source	Reporting Unit	Population	Description
				measurement year age 18 years and older
Children Get Immunization Shot	DOH	Percentage	All	The percentage of children 2 years of age who had one dose of diphtheria, tetanus and acellular pertussis (IPV); one measles, mumps and rubella (MMR); one haemophilus influenza type B (HiB); one chicken pox (VZV); four pneumococcal (Pneumo); one hepatitis (Hep A); two or three rotavirus (flu) vaccines by their second birthday.
Teenagers Get Immunization Shots	DOH	Percentage	All	The percentage of children 13 years of age who had one dose of meningococcal conjugate vaccine, one tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccine and three doses of the human papillomavirus (HPV) vaccine by their 13th birthday, Combination Rate 2; (2) HPV for Female Adolescents; and (3) HPV for Male Adolescents
Girls Get HPV Shot to Prevent Infection and Disease	DOH	Percentage	All	The percentage of children 13 years of age who had three doses of the human papillomavirus (HPV) vaccine by their 13th birthday, HPV for Female Adolescents
Boys Get HPV Shot to Prevent Infection and Disease	DOH	Percentage	All	The percentage of children 13 years of age who had three doses of the human papillomavirus (HPV) vaccine by their 13th birthday, HPV for Male Adolescents
Children Get Flu Shot	DOH	Percentage	All	Percentage of patients aged 6 months and older seen for a visit and who received an influenza immunization OR who reported previous receipt of an influenza immunization, ages 6 months - 17 years
Care to Prevent Cavities	HCA	Percentage	Medicaid	Total number of patients (Age < 6), who received a Fluoride Varnish application during a routine preventive health visit.
Testing Hearing Loss in Infants	DOH	Percentage	All	The percentage of newborns who did not pass hearing screening and have an audiological evaluation no later than 3 months of age.

Measure Name	Source	Reporting Unit	Population	Description
Checking for Breast Cancer	Claims	Percentage	Medicaid	The percentage of women, ages 50–74 years, who had a mammogram to screen for breast cancer.
Checking for Cervical Cancer	Claims	Percentage	Medicaid	The percentage of women, ages 21–64 years, who were screened for cervical cancer using either of two methods defined by the measure (interval dependent upon screening method).
Adults Get Flu Shot	DOH	Percentage	All	Percentage of patients aged 6 months and older seen for a visit and who received an influenza immunization OR who reported previous receipt of an influenza immunization, ages 18 and older
Check for Chlamydia Infection in Younger Women	Claims	Percentage	Medicaid	The percentage of women, ages 16–24 years, who were identified as sexually active and who had at least one test for chlamydia during the measurement year.
Older Adults Get Pneumonia Vaccine Shot	DOH	Percentage	All	The percentage of patients 65 years of age and older who have ever received a pneumococcal vaccine.

WA-APCD INSURANCE MARKET SECTOR QUALITY MEASURES

Measure Name	Source	Reporting Unit	Population	Description
Yearly Healthcare Cost per Medicaid Enrollee	HCA	Per Enrollee Per Year	Medicaid	Total Medicaid spending in the calendar year divided by the total number of Medicaid beneficiaries in the calendar year.
Yearly Healthcare Cost per Public Employee Program Enrollee (PEBB)	HCA	Per Enrollee Per Year	PEBB	Total State spending for public employees and dependents in the calendar year divided by the total number of beneficiaries in the calendar year.
Emergency Room Visits Per 1,000 Persons	Claims	Visits Per 1,000 Persons	Commercial Exchange Medicaid	Number of emergency department visits per 1,000 population and is calculated in member months for Medicaid and member years for commercial/Exchange.

Measure Name	Source	Reporting Unit	Population	Description
Avoid Return to Hospital	Claims	Ratio Observed to Expected	Commercial	For members 18 years of age and older, the number of acute inpatient stays that were followed by an unplanned acute readmission for any diagnosis within 30 days and the predicted probability of an acute readmission.
Avoid Return to Hospital for Mental Health Condition	DSHS	Ratio Observed to Expected	Medicaid	For members 18 years of age and older, the number of acute inpatient psychiatric stays that were followed by an acute readmission for a psychiatric diagnosis within 30 days. Report for Medicaid only.
Controlling High Blood Pressure	NCQA	Percentage	Commercial Exchange Medicaid	The percentage of patients 18 to 85 years of age who had a diagnosis of hypertension (HTN) and whose blood pressure (BP) was adequately controlled (<140/90) during the measurement year.
Blood Sugar in Good Control - Diabetes	NCQA	Percentage	Commercial Medicaid	The percentage of patients 18-75 years of age with diabetes (type 1 and type 2) whose most recent HbA1c level during the measurement year was greater than 9.0% (poor control) or was missing a result, or if an HbA1c test was not done during the measurement year.
Blood Pressure in Good Control - Diabetes	NCQA	Percentage	Commercial Medicaid	The percentage of patients 18-75 years of age with diabetes (type 1 and type 2) whose most recent blood pressure level taken during the measurement year is <140/90 mm Hg.
Testing for Cause of Low Back Pain	Claims	Percentage	Commercial Exchange Medicaid	The percentage of members, ages 18-50 years, with a primary diagnosis of low back pain who did not have an imaging study (i.e., plain X-ray, MRI, or CT scan) within 28 days of the diagnosis. (Note that a higher score indicates better performance.)
Avoid Wrong Medication for Bronchitis	Claims	Percentage	Commercial Exchange Medicaid	The percentage of adults, ages 18-64 years, with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription.
Checking Patients Who Need Ongoing Medications	Claims	Percentage	Commercial Exchange Medicaid	The percentage of patients, ages 18 years and older, who received at least 180 treatment days of ambulatory medication therapy for ACE/ARB during the measurement year and had at least one therapeutic monitoring event for the therapeutic agent in the measurement year.
Treating Children with Throat Infections	Claims	Percentage	Commercial Exchange Medicaid	The percentage of children, ages 3-18 years, who were diagnosed with pharyngitis, dispensed an antibiotic, and received a group A streptococcus (strep) test for the episode. A higher rate represents better performance (i.e., appropriate testing).

Measure Name	Source	Reporting Unit	Population	Description
Avoid Unnecessary Emergency Visits	Claims	Rate per 1,000 Member Years	Commercial Exchange Medicaid	The rate of patient ED visits for conditions that could have been more appropriately managed by or referred to a primary care provider.
Blood Sugar Tested	Claims	Percentage	Commercial Exchange Medicaid	The percentage of members, ages 18–75 years, with diabetes (Type 1 and/or Type 2) who had an HbA1c test during the measurement year.
Check for Kidney Disease	Claims	Percentage	Commercial Exchange Medicaid	The percentage of members, ages 18–75 years, with diabetes (Type 1 and/or Type 2) who received a nephropathy screening or monitoring test or had evidence of nephropathy during the measurement year or the year prior.
Eye Exam to Check for Vision Loss	Claims	Percentage	Commercial Exchange Medicaid	The percentage of members, ages 18–75 years, with diabetes (Type 1 and/or Type 2) who had an eye exam during the measurement year or the year prior.
Getting Medications to Control Blood Sugar	Claims	Percentage	Commercial Exchange Medicaid	The percentage of patients, ages 18 years and older, who met the proportion of days covered (PDC) threshold of 80% for diabetes roll-up medications.
Getting Anti-depressant Medications	Claims	Percentage	Commercial Exchange Medicaid	The percentage of members 18 years and older who were treated with antidepressant medication, had a diagnosis of major depression and who remained on an antidepressant medication treatment: effective acute phase treatment rate
Getting Care for Attention Deficit/Hyperactivity Disorder	Claims	Percentage	Commercial Exchange Medicaid	The percentage of children newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed. Continuation and Maintenance Phase rate
Follow-Up Visit After Hospital Stay for Mental Health	Claims	Percentage	Commercial Exchange Medicaid	The percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness diagnoses and who had a follow-up visit with a mental health practitioner within 30 days of discharge
Testing for Lung Disease	Claims	Percentage	Commercial Exchange Medicaid	The percentage of members 40 years of age and older with a new diagnosis of COPD or newly active COPD, who received appropriate spirometry testing to confirm the diagnosis.

Measure Name	Source	Reporting Unit	Population	Description
Getting Medications for Asthma	Claims	Percentage	Commercial Exchange Medicaid	The percentage of members 5-85 years of age who were identified as having persistent asthma and were dispensed appropriate medications that they remained on for at least 75% of their treatment period. (For Medicaid, this measure is reported for people ages 5-64.)
Fewer Hospital Stays Due to Good Lung Care	AHRQ	Rate per 1,000 Member Years	Commercial Exchange Medicaid	The rate of adults, ages 40 years and older, who had an admission for a principal diagnosis of chronic obstructive pulmonary disease (COPD) or asthma.
Getting Medications to Lower Blood Pressure	Claims	Percentage	Commercial Exchange Medicaid	The percentage of patients, ages 18 years and older, who met the proportion of days covered (PDC) threshold of 80% for Renin Angiotensin System Antagonists.
Getting Medications to Lower Cholesterol	Claims	Percentage	Commercial Exchange Medicaid	The percentage of patients, ages 18 years and older, who met the proportion of days covered (PDC) threshold of 80% for statins
Infants Get Regular Check-ups	Claims	Percentage	Commercial Exchange Medicaid	The percentage of members who turned 15 months old during the measurement year and who had six or more well-child visits with a PCP during their first 15 months of life.
Children Get Regular Check-ups	Claims	Percentage	Commercial Exchange Medicaid	The percentage of members, ages 3-6 years, who had one or more well-child visits with a PCP during the measurement year.
Having a Primary Care Visit Yearly	Claims	Percentage	Commercial Exchange Medicaid	The percentage of members 12 months - 19 years of age who had a visit with a PCP, all children ages.
Getting Substance Abuse Services	DSHS	Percentage	Medicaid	The percentage of members with a substance use disorder treatment need who received a substance use disorder treatment in the measurement year, age 12-17 years
Getting Mental Health Services	DSHS	Percentage	Medicaid	The percentage of members with a mental health service need who received mental health services in the measurement year age 6-17 years
Getting Mental Health Services	DSHS	Percentage	Commercial	The percentage of members with a mental health service need who received mental health services in the measurement year age 6-17 years

Measure Name	Source	Reporting Unit	Population	Description
Having a Doctor/Office Visit Yearly	Claims	Percentage	Commercial Exchange Medicaid	The percentage of members 20 years and older who had an ambulatory or preventive care visit, all adult ages.
Getting Substance Abuse Services	DSHS	Percentage	Medicaid	The percentage of members with a substance use disorder treatment need who received a substance use disorder treatment in the measurement year, age 18 years and older
Getting Mental Health Services	DSHS	Percentage	Medicaid	The percentage of members with a mental health service need who received mental health services in the measurement year age 18 years and older
Getting Mental Health Services	DSHS	Percentage	Commercial	The percentage of members with a mental health service need who received mental health services in the measurement year age 18 years and older
Checking if Weight is a Health Problem – Children	NCQA	Percentage	Commercial Medicaid	The percentage of members 3-17 years of age who had an outpatient visit with a PCP or OB/GYN and who had evidence of the following during the measurement year: (1) BMI percentile documentation; (2) counseling for nutrition; and (3) counseling for physical activity, children all ages.
Care to Prevent Cavities	HCA	Percentage	Medicaid	Total number of patients (Age < 6), who received a Fluoride Varnish application during a routine preventive health visit.
Checking for Breast Cancer	Claims	Percentage	Commercial Exchange Medicaid	The percentage of women, ages 50–74 years, who had a mammogram to screen for breast cancer.
Checking for Cervical Cancer	Claims	Percentage	Commercial Exchange Medicaid	The percentage of women, ages 21–64 years, who were screened for cervical cancer using either of two methods defined by the measure (interval dependent upon screening method).
Checking for Colorectal Cancer	NCQA	Percentage	Commercial	The percentage of members 50-75 years of age who had appropriate screening for colorectal cancer. (interval dependent upon screening method).
Checking if Weight is a Health Problem - Adults	NCQA	Percentage	Commercial Medicaid	The percentage of members 18-74 years of age who had an outpatient visit and whose body mass index (BMI) was documented during the measurement year or the year prior to the measurement year.

Measure Name	Source	Reporting Unit	Population	Description
Check for Chlamydia Infection in Younger Women	Claims	Percentage	Commercial Exchange Medicaid	The percentage of women, ages 16–24 years, who were identified as sexually active and who had at least one test for chlamydia during the measurement year.